



EVOLUZIONE 2017

7° Congresso della Società Italiana di Biologia Evoluzionistica

Abstracts book

Rome, 28-31 August 2017

Department of Sciences, Roma Tre University, viale G.Marconi 446

Organising Committee

GABRIELE GENTILE (Coord.)

Roma Tor Vergata

GIULIANA ALLEGROCCI

Roma Tor Vergata

MARCO ALBERTO BOLOGNA

Roma Tre

MONICA CAROSI

Roma Tre

DONATELLA CESARONI

Roma Tor Vergata

ALESSIO DE BIASE

Roma Sapienza

ANDREA DI GIULIO

Roma Tre

EMILIANO MANCINI

Roma Tre

MARCO OLIVERIO

Roma Sapienza

LEONARDO VIGNOLI

Roma Tre

Organising Secretariat

CRISTINA COMPAGNO

Management of Tourism and

Biodiversity - Elly Travel srl

info@sibe2017.uniroma2.it

Scientific Committee

MARCO PASSAMONTI (Coord.)

Bologna

GIULIANA ALLEGROCCI

Roma Tor Vergata

GIORGIO BERTORELLE

Ferrara

MAURIZIO CASIRAGHI

Milano Bicocca

PAOLO CIUCCI

Roma Sapienza

GIUSEPPE FUSCO

Padova

GABRIELE GENTILE

Roma Tor Vergata

SILVIA GHIROTTI

Ferrara

ALESSANDRO MINELLI

Padova

TELMO PIEVANI

Padova

ANDREA PILASTRO

Padova

OMAR ROTA STABELLI

Edmund Mach Foundation (TN)

Sponsors



Dating the impossible: the origin and divergence of the largest infection on Earth

Francesco Drago¹, Matthias Scholz², Nicola Segata³, Omar Rota-Stabelli²

¹ Università degli Studi di Padova,

² Fondazione Edmund Mach,

³ CIBIO, Università degli Studi di Trento

Wolbachia are common intracellular bacteria of many arthropod and nematode species; they are responsible for likely the most widespread and evolutionary significant infection on Earth. *Wolbachia* are generally acquired by maternal transmission, but they often move across species boundaries, impairing the use of host divergences as independent calibration to infer their evolutionary history. Here we present a first systematic effort to estimate the origin and the divergence of the *Wolbachia* infection by coupling the molecular clock analysis of a genomic and an MLST data sets with three types of disputed calibration priors: a root prior for splits within Proteobacteria, the recently proposed *Nomada* host-symbiont co-divergence, and an adjusted mutation rate gathered from *Drosophila melanogaster* and *Nasonia* species. We first perform a hierarchical based model comparison on the MLST data set to select the best clock, replacement, and tree priors; we then used the best prior combination to calibrate both the genomic and the MLST data sets using different combinations of calibrations. Although divergence estimates are characterized by a high degree of uncertainty and strongly depend on the use of certain calibrations, most results point towards a pre-Cambrian origin of *Wolbachia* (its split from *Ehrlichia*). Diversification of *Wolbachia* supergroups is more recent, from the late Paleozoic to the Mesozoic: this is compatible with a long-fuse model of evolution, and suggests a co-radiation with holometabolan insects. Our results provide a comprehensive first effort to date *Wolbachia* evolution: results are model, prior, and data set dependent and reveal the difficulties embedded in the molecular dating of such elusive bacteria characterized by the absence of reliable calibration priors and a clear evolutionary history.